

Radio Properties of EROs in the Phoenix Deep Survey

J. Afonso¹, A. M. Hopkins², M. Sullivan³, B. Mobasher⁴, A. Georgakakis⁵, and
L. E. Cram⁶

¹ CAAUL, Observatory of Lisbon, Tapada da Ajuda, 1349-018 Lisbon, Portugal

² Dept. of Physics & Astronomy, University of Pittsburgh, 3941 O'Hara St,
Pittsburgh, PA 15260, USA

³ Dept. of Astronomy and Astrophysics, University of Toronto, 60 St. George St,
Toronto, Ontario M5S 3H8, Canada

⁴ ESA, STScI, 3700 San Martin Drive, Baltimore, MD 21218, USA

⁵ National Athens Observatory, I.Metaxa & Vas.Pavlou str., Athens 15236, Greece

⁶ Australian Research Council, GPO Box 2702, Canberra, ACT 2601, Australia

Abstract. Insensitive to dust obscuration, radio wavelengths are ideal to study star-forming galaxies free of dust induced biases. Using data from the Phoenix Deep Survey, we have identified a sample of star-forming extremely red objects (EROs). Stacking of the radio images of the radio-undetected star-forming EROs revealed a significant radio detection. Using the expected median redshift, we estimate an average star-formation rate of $61 \text{ M}_{\odot} \text{ yr}^{-1}$ for these galaxies.

1 Introduction

Identifying and studying dusty star-forming galaxies is of vital importance to understand galaxy evolution. Besides looking directly for the emission from dust, these objects can also be identified by their extreme optical to near-infrared (NIR) colours, forming a significant fraction of the so-called Extremely Red Objects (EROs) [3]. Radio wavelengths, unaffected by dust obscuration and being a measure of star formation rates in star-forming galaxies, can give an important insight to the characteristics of these dusty objects [1]. Performing an ERO search in an ultra-deep radio field can thus provide an important step in understanding some of the more obscured galaxies.

2 Star-forming EROs in the PDS

The Phoenix Deep Survey is a multiwavelength survey based on deep radio (1.4 GHz) observations performed with the ATCA. With an area of over 4.5 square degree, and reaching flux densities of $60 \mu\text{Jy}$, the PDS is one of the deepest wide angle radio surveys already made [4]. Follow-up optical (*UBVRI*) data has recently been obtained, using the Wide-Field Imagers at the AAT, CTIO and ESO, reaching ~ 25 mag in each band. Near-infrared (K_s band) observations were made on the central 200 square arcmin of the PDS, using SOFI at the NTT (ESO) to a limiting magnitude of $K_s \sim 20$ [5]. Based on $3''$ aperture photometry,




Fig. 1. Classification of EROs using the RIK_s colour-colour plot (left) and the stacked radio image of the radio-undetected star-forming EROs, revealing a $8.7 \mu\text{Jy}$ detection.

a sample of 432 EROs ($R - K_s > 5$), were selected. Eighteen were directly identified in the radio catalogue, with radio fluxes between 65 and $991 \mu\text{Jy}$. The ERO population was separated into ellipticals and star-forming galaxies using the RIK_s magnitudes, following the recent work by Bergstrom et al. [2]. This method seems to perform well for EROs with redshifts $z < 1.4$ [2], which seems appropriate for this survey (a spectroscopic analysis of an ERO sample with similar K_s magnitude limits revealed a median redshift of $z \sim 1.2$, with a very small number of EROs having $z > 1.4$ [3]). A total of 214 objects fall in the star-forming region of the RIK_s colour-colour plot (Figure 1). After removing all ($> 3\sigma$) radio detections, the radio images at the locations of the star-forming EROs were stacked. A significant (6.7σ) detection of $8.7 \mu\text{Jy}$ was revealed (Figure 1). Assuming a median redshift for the sample of $z \sim 1.2$, and assuming the radio emission is solely due to star-formation, this implies an average star-formation rate of $61 \text{ M}_\odot \text{ yr}^{-1}$.

Improvement on the photometric analysis, by using the full photometric dataset to estimate photometric redshifts and classifications, is currently underway, and will result in a precise determination of the contribution of star-forming EROs to the star-formation rate in the Universe.

AMH acknowledges support provided by NASA through Hubble Fellowship grant HST-HF-01140.01-A awarded by STScI. JA acknowledges support from the Science and Technology Foundation (FCT, Portugal) through the fellowship BPD-5535-2001 and the research grant POCTI-FNU-43805-2001.

References

1. J. Afonso, B. Mobasher, B. Chan, L. Cram: ApJ, **559**, L101 (2001)
2. S. Bergström, T. Wiklind: A&A (2003) (in press; astro-ph/0310628)
3. A. Cimatti, et al.: A&A (2003) (in press; astro-ph/0310742)
4. A. M. Hopkins, J. Afonso, B. Chan, L. E. Cram, A. Georgakakis, B. Mobasher: AJ, **125**, 465 (2003)
5. M. Sullivan, A. M. Hopkins, J. Afonso, A. Georgakakis, B. Chan, L. E. Cram, B. Mobasher: ApJS, submitted.